

Free Online
Access for
Foreign Institutions


ISSN 0972-0871

This title
is indexed
in Scopus



Far East Journal of Mathematical Sciences (FJMS)

Volume 69, Number 2
October 16-31, 2012



Pushpa Publishing House

Vijaya Niwas, 198 Mumfordganj
Allahabad 211002, INDIA

<http://pphmj.com/journals/fjms.htm>
fjms@pphmj.com & arun@pphmj.com



ON SUPER EDGE ANTI MAGIC TOTAL LABELING FOR t -JOINT COPIES OF WHEEL

I W. Sudarsana¹, A. Hendra¹, Adiwijaya² and D. Y. Setyawan³

¹Combinatorial and Applied Mathematics Research Group
Tadulako University
Jalan Sukarno-Hatta Km. 9 Palu 94118, Indonesia
e-mail: sudarsanaiwayan@yahoo.co.id
beyourselft@yahoo.co.id

²Algorithm and Computation Research Group
Institut Teknologi Telkom
Jalan Telekomunikasi No. 1
Terusan Buah Batu Bandung 40257, Indonesia
e-mail: adiwijaya007@yahoo.co.id

³Institut Bisnis dan Informatika Darmajaya
Jalan Z. A. Pagar Alam No. 93 Bandar Lampung 35141
Indonesia
e-mail: dodi.yudo@darmajaya.ac.id

Abstract

A (p, q) -graph G is called (a, d) -edge anti magic total, (a, d) -EAMT, if there exist integers $a > 0$, $d \geq 0$ and a bijection $\lambda : V \cup E \rightarrow \{1, 2, \dots, p + q\}$ such that $W = \{w(xy) : xy \in E\} = \{a, a + d, \dots, a$

© 2012 Pushpa Publishing House

2010 Mathematics Subject Classification: 05C78.

Keywords and phrases: (a, d) -EAMT labeling, (a, d) -SEAMT labeling, dual labeling.

Received April 13, 2012

276 I W. Sudarsana, A. Hendra, Adiwijaya and D. Y. Setyawan
 $+ (q - 1)d$, where $w(xy) = \lambda(x) + \lambda(y) + \lambda(xy)$ is the edge weight of xy . An (a, d) -EAMT labeling λ of G is called super, (a, d) -SEAMT if $\lambda(V) = \{1, 2, \dots, p\}$. In this paper, we show that t -joint copies of wheel have a $((2n + 2)t + 2, 1)$ -SEAMT labeling for even $n \geq 4$, and $t \geq 2$.

1. Introduction

We consider finite undirected graphs without loops and multiple edges. The notation $V(G)$ and $E(G)$ stand for the vertex set and edge set of graph G , respectively. Let $e = \{u, v\}$ (in short, $e = uv$) denote an edge connecting vertices u and v in G . A graph P_n denotes a path on n vertices. For other standard terminologies and notation for graph theoretic ideas, we follow the book of [5].

We denote by (p, q) -graph G a graph with p vertices and q edges. A (p, q) -graph G is called (a, d) -edge anti magic total, (a, d) -EAMT, if there exist integers $a > 0$, $d \geq 0$ and a bijection $\lambda: V \cup E \rightarrow \{1, 2, \dots, p + q\}$ such that the set of edge-weights is

$$W = \{w(xy) : xy \in E\} = \{a, a + d, \dots, a + (q - 1)d\},$$

where $w(xy) = \lambda(x) + \lambda(y) + \lambda(xy)$. We shall follow [7] to call $w(xy) = \lambda(x) + \lambda(y) + \lambda(xy)$ the *edge-weight* of xy , and W the *set of edge-weights* of the graph G . In particular, an (a, d) -EAMT labeling λ of a (p, q) -graph G is *super* if $\lambda(V) = \{1, 2, \dots, p\}$. For the rest of paper, we will denote super (a, d) -EAMT of G by (a, d) -SEAMT.

For any (a, d) -SEAMT labeling on a (p, q) -graph G , the maximum edge-weight is no more than $p + (p - 1) + (p + q)$. Thus, $a + (q - 1)d \leq 3p + q - 1$. Similarly, the minimum possible edge-weight is at least $1 + 2 + p + 1$. This implies that $a \geq p + 4$. Therefore, we have

$$d \leq \frac{2p + q - 5}{q - 1}. \quad (1)$$

In general, for any (a, d) -EAMT labeling on a (p, q) -graph G , the maximum edge-weight is no more than $(p + q - 2) + (p + q - 1) + (p + q)$. Thus, $a + (q - 1)d \leq 3p + 3q - 3$. Similarly, the minimum possible edge-weight is at least $1 + 2 + 3$. Consequently, $a \geq 6$. So, we have

$$d \leq \frac{3p + 3q - 9}{q - 1}. \quad (2)$$

A number of classification studies on (a, d) -SEAMT (resp. (a, d) -EAMT) for connected graphs has been extensively investigated. For instances, in [2], Bača et al. showed that wheel W_n has an (a, d) -SEAMT labeling if and only if $d = 1$ and $n \equiv 1 \pmod{4}$; Fan F_n has an (a, d) -SEAMT if $2 \leq n \leq 6$ and $d \in \{0, 1, 2\}$. Ngurah and Baskoro [6] proved that for every Petersen graph $P(n, m)$, $n \geq 3$, $1 \leq m \leq \frac{n}{2}$, has a $(4n + 2, 1)$ -SEAMT labeling. More results concerning SEAMT graphs can be seen in [1] and survey paper by Gallian [4]. People also consider how to construct a new (bigger) (a, d) -SEAMT graph from some known (smaller) (a, d) -SEAMT graphs. These constructions are proposed by inserting some new pendant edges and points, see for instance [3, 7, 8, 10, 11]. However, the (a, d) -SEAMT labeling for t -joint copies of wheel, $W_{(t, n)}$, is still open.

In this paper, we show that $W_{(t, n)}$ has a $((2n + 2)t + 2, 1)$ -SEAMT labeling for even $n \geq 4$, and $t \geq 2$.

2. Preliminary Theorems

The properties of (a, d) -SEAMT labeling of graph proposed in the following theorems will be useful in the next section. Given any (a, d) -EAMT labeling λ on a (p, q) -graph G . Then its *dual* labeling λ' can be defined [12] by

$$\lambda'(x) = p + q + 1 - \lambda(x) \text{ for any vertex } x, \text{ and}$$

$$\lambda'(xy) = p + q + 1 - \lambda(xy) \text{ for any edge } xy.$$

By using this duality, Wallis et al. [12] established the following theorem.

Theorem A. (Wallis et al. [12]). *If a (p, q) -graph G has an (a, d) -EAMT labeling, then G has an (a', d) -EAMT labeling as its dual with $a' = 3p + 3q + 3 - a - (q - 1)d$.*

Theorem B. (Sudarsana et al. [9]). *Let λ be an (a, d) -SEAMT labeling of a (p, q) -graph G . Then the labeling λ' defined:*

$$\lambda'(x) = p + 1 - \lambda_1(x), \forall x \in V \text{ and}$$

$$\lambda'(xy) = 2p + q + 1 - \lambda_1(xy), \forall xy \in E$$

is an (a', d) -SEAMT labeling of G with $a' = 4p + q + 3 - a - (q - 1)d$.

The labeling λ' is called a *dual (a, d) -SEAMT labeling* of λ on G . Moreover, the dual labeling is called *selfdual*, if $a = a'$.

3. The Main Result

Let W_n be a wheel on $n + 1$ vertices with v_0 as the *hub* and cycle $v_1v_2v_3 \cdots v_nv_1$ as the *rim*. The *t-joint copies of wheel*, denoted by $W_{(t,n)}$, is a graph obtained by taking t copies of W_n and joining every hub of the wheels by an edge such that forming a path P_t . We provide an example of $W_{(4,6)}$ in Figure 1:




Figure 1. The 4-joint copies of $W_6, W_{(4,6)}$.

In general, we denote that $V(W_{t,n}) = \{v_{i,j} : 1 \leq j \leq t, 0 \leq i \leq n\}$ and $E(W_{t,n}) = \{e_{i,j}, e_{n,j}, e_{i,j}^o, e_j^o\}$, where

$$e_{i,j} = \begin{cases} v_{i,j}v_{i+1,j}, & 1 \leq i \leq n-1, \quad 1 \leq j \leq t; \\ v_{n,j}v_{1,j}, & i = n, \quad 1 \leq j \leq t, \end{cases}$$

$$e_{i,j}^o = v_{0,j}v_{i,j}, \quad 1 \leq j \leq t, \quad 1 \leq i \leq n,$$

$$e_j^o = v_{0,j}v_{0,j+1}, \quad 1 \leq j \leq t-1.$$

By (1) and (2), we have: for every even $n \geq 4$, and $t \geq 2$ is an integer, there is no (a, d) -SEAMT labeling of $W_{(t,n)}$ with $d \geq 3$; and there is no (a, d) -EAMT labeling of $W_{(t,n)}$ with $d \geq 5$.

The following theorem deals with $((2n+2)t+2, 1)$ -SEAMT labeling of $W_{(t,n)}$.

Theorem 1. *Let $n \geq 4$ and $t \geq 2$ be integers, and n is even. Then the graph $W_{(t,n)}$ has a $((2n+2)t+2, 1)$ -SEAMT labeling. This type of labeling is selfdual.*

Proof. Label the vertices and edges of $W_{(t,n)}$ in the following way:

$$\lambda(v_{i,j}) = \begin{cases} (i-1)t+j, & 1 \leq i \leq \frac{n}{2}, \quad 1 \leq j \leq t; \\ \frac{nt}{2}+j, & i=0, \quad 1 \leq j \leq t; \\ it+j, & \frac{(n+2)}{2} \leq i \leq n, \quad 1 \leq j \leq t, \end{cases}$$

$$\lambda(e_{i,j}) = \begin{cases} (2n+2-i)t+1-j, & 1 \leq i \leq \frac{n}{2}, \quad 1 \leq j \leq t; \\ (2n+1-i)t+1-j, & \frac{(n+2)}{2} \leq i \leq n-1, \quad 1 \leq j \leq t; \\ \left(\frac{3n+2}{2}\right)t+1-j, & i=n, \quad 1 \leq j \leq t, \end{cases}$$

$$\lambda(e_{i,j}^o) = \begin{cases} (3n+3-2i)t+1-j, & 1 \leq i \leq \frac{n}{2}, \quad 1 \leq j \leq t; \\ (4n+2-2i)t+1-j, & \frac{(n+2)}{2} \leq i \leq n, \quad 1 \leq j \leq t, \end{cases}$$

$$\lambda(e_j^o) = (3n+2)t-j, \quad 1 \leq j \leq t.$$

Now, we construct the set of edge-weights W in the following way:

$$W = \{w(e_{i,j}), w(e_{i,j}^o), w(e_j^o)\},$$

where

$$w(e_{i,j}) = \begin{cases} \lambda(v_{i,j}) + \lambda(e_{i,j}) + \lambda(v_{i+1,j}), & 1 \leq i \leq n-1, \quad 1 \leq j \leq t; \\ \lambda(v_{n,j}) + \lambda(e_{n,j}) + \lambda(v_{1,j}), & i = n, \quad 1 \leq j \leq t, \end{cases}$$

$$w(e_{i,j}^o) = \lambda(v_{0,j}) + \lambda(e_{i,j}^o) + \lambda(v_{i,j}), \quad 1 \leq i \leq n, \quad 1 \leq j \leq t;$$

$$w(e_j^o) = \lambda(v_{0,j}) + \lambda(e_j^o) + \lambda(v_{0,j+1}), \quad 1 \leq j \leq t-1.$$

Therefore, we obtain that the edge-weights are

$$w(e_{i,j}) = \begin{cases} (2n+1+i)t+1+j, & 1 \leq i \leq \frac{n}{2}-1, \quad 1 \leq j \leq t; \\ \left(\frac{5n+4}{2}\right)t+1+j, & i = \frac{n}{2}, \quad 1 \leq j \leq t; \\ (2n+2+i)t+1+j, & \frac{(n+2)}{2} \leq i \leq n-1, \quad 1 \leq j \leq t; \\ \left(\frac{5n+2}{2}\right)t+1+j, & i = n, \quad 1 \leq j \leq t, \end{cases}$$

$$w(e_{i,j}^o) = \begin{cases} \left(\frac{7n+5}{2}-i\right)t+1+j, & 1 \leq i \leq \frac{n}{2}, \quad 1 \leq j \leq t; \\ \left(\frac{9n+4}{2}-i\right)t+1+j, & \left(\frac{(n+2)}{2}\right) \leq i \leq n, \quad 1 \leq j \leq t, \end{cases}$$

$$w(e_j^o) = (4n+2)t+1+j, \quad 1 \leq j \leq t-1.$$

Finally, we have that the set of edge-weights is

$$W = \{(2n + 2)t + 2, (2n + 2)t + 3, \dots, (4n + 3)t - 1, (4n + 3)t\}.$$

We can see that the set W consists of the consecutive integers starting from $a = (2n + 2)t + 2$. By Theorem B, it can be verified that this labeling is selfdual. The proof is now complete. \square




Figure 2. A $(44, 1)$ -SEAMT labeling of $W_{(3,6)}$ obtained from Theorem 1.

By Theorem A, we have the following corollary.

Corollary 1. For every even $n \geq 4$ and $t \geq 2$ is an integer, the graph $W_{(t,n)}$ has an $((5n + 3)t, 1)$ -EAMT labeling.

To conclude this paper, let us present two open problems to work on. Construct, if there exists




Figure 3. An $(99, 1)$ -EAMT labeling of $W_{(3,6)}$ obtained from Theorems 1 and A.

1. An (a, d) -SEAMT labeling of $W_{(t,n)}$ with $d \in \{0, 2\}$, for even $n \geq 4$, and $t \geq 2$.

2. An (a, d) -SEAMT labeling of $W_{(t,n)}$ with $d \in \{0, 1, 2\}$, for odd $n \geq 5$, and $t \geq 2$.

Acknowledgement

The first and second authors gratefully acknowledge to the Directorate General of Higher Education of Indonesia (DIKTI), State Ministry of Education and Culture for financial support under Penelitian Fundamental: 125.C/un 28.2/PL/2012.

References

- [1] M. Bača, Y. Lin, M. Miller and R. Simanjuntak, New constructions of magic and antimagic graph labelings, *Utilitas Math.* 60 (2001), 229-239.
- [2] M. Bača, Y. Lin, M. Miller and M. Z. Youssef, Edge-antimagic graphs, *Discrete Math.* 307 (2007), 1232-1244.
- [3] E. T. Baskoro, I W. Sudarsana and Y. M. Cholily, How to construct new super edge-magic graphs from some old ones, *J. Indones. Math. Soc.* 11(2) (2005), 156-162.
- [4] J. A. Gallian, A dynamic survey of graph labellings, *Electron. J. Combin.* 18 (2011), #DS6.
- [5] N. Hartsfield and G. Ringel, *Pearls in Graph Theory*, Academic Press, San Diego, 1994.
- [6] A. A. G. Ngurah and E. T. Baskoro, On magic and antimagic total labeling of generalized Petersen graph, *Utilitas Math.* 63 (2003), 97-107.
- [7] Kiki A. Sugeng and Mirka Miller, Relationship between adjacency matrices and super (a, d) -edge-antimagic-total labeling of graphs, *J. Combin. Math. Combin. Comput.* 55 (2005), 71-82.
- [8] I W. Sudarsana, E. T. Baskoro, D. Izmailusa and H. Assiyatun, Creating new super edge-magic total labelings from old ones, *J. Combin. Math. Combin. Comput.* 55 (2005), 83-90.
- [9] I W. Sudarsana, E. T. Baskoro, D. Izmailusa and H. Assiyatun, On super (a, d) -edge antimagic total labeling of disconnected graphs, *J. Combin. Math. Combin. Comput.* 55 (2005), 149-158.

- [10] I. W. Sudarsana, E. T. Baskoro, S. Uttungadewa and D. Izmailusa, An expansion technique on super edge-magic total graphs, *Ars Combin.* 91 (2009), 231-241.
- [11] I. W. Sudarsana, E. T. Baskoro, S. Uttungadewa and D. Izmailusa, Expansion techniques on the super edge anti magic total graphs, *J. Combin. Math. Combin. Comput.* 71 (2009), 189-199.
- [12] W. D. Wallis, E. T. Baskoro, M. Miller and Slamin, Edge-magic total labellings, *Australian J. Combin.* 22 (2000), 177-190.