

JCMCC

THE JOURNAL

of

COMBINATORIAL MATHEMATICS

and

COMBINATORIAL COMPUTING

THE JOURNAL OF COMBINATORIAL MATHEMATICS
AND COMBINATORIAL COMPUTING

ISSN 0835-3026

EDITOR-IN-CHIEF

R.E.L. Aldred
University of Otago
Box 56, Dunedin, New Zealand
raldred@maths.otago.ac.nz

EDITORIAL BOARD

A.H. Baartmans
Michigan Technological University
Houghton, MI, USA 49931-1295
baartman@math.mtu.edu

E.J. Billington
University of Queensland
Queensland 4072, Australia
ejb@maths.uq.edu.au

L. Caccetta
Curtin University of Technology
Box U1987, Perth, WA 6845, Australia
caccetta@cs.curtin.edu.au

L.J. Cummings
University of Waterloo
Waterloo, ON, Canada N2L 3G1
ljcumming@math.uwaterloo.ca

R. Fuji-Hara
University of Tsukuba
1-1-1 Tennodai, Tsukuba 305-8573, Japan
fujihara@sk.tsukuba.ac.jp

R.J. Gould
Emory University
Atlanta, GA USA 30322-2390
rg@mathcs.emory.edu

T.W. Haynes
East Tennessee State University
Johnson City, TN, USA 37614-0663
haynes@mail.etsu.edu

A.J.W. Hilton
University of Reading
Whiteknights, Reading RG6 2AX, Great Britain
a.j.w.hilton@reading.ac.uk

P. Horak
University of Washington, Tacoma
Tacoma, WA USA 98402
horak@u.washington.edu

C.S. Iliopoulos
King's College London
Strand, London WC2R 2LS, Great Britain
csi@dcs.kcl.ac.uk

M. Jimbo
Keio University, Hiyoshi Campus
3-14-1 Hiyoshi, Yokohama 223-8522, Japan
jimbo@math.keio.ac.jp

D. Jungnickel
University Augsburg
D-86135 Augsburg, Germany
jungnickel@math.uni-augsburg.de

D. Kreher
Michigan Technological University
Houghton, MI, USA 49931-1295
kreher@math.mtu.edu

J. Leung
New Jersey Institute of Technology
Newark, NJ, USA 07102
leung@homer.njit.edu

C.M. Mynhardt
University of Victoria
Victoria, BC Canada V8W 2Y2
mynhardt@math.uvic.ca

R. Rees
Memorial University of Newfoundland
St John's, NL Canada A1C 5S7
rolf@math.mun.ca

R.W. Robinson
University of Georgia
Athens, GA, USA 30602-7404
rwr@cs.uga.edu

C.A. Rodger
Auburn University
Auburn, AL, USA 36849-5307
rodger1@mail.auburn.edu

R.G. Stanton
University of Manitoba
Winnipeg, MB, Canada R3T 2N2
stanton@cc.umanitoba.ca

D.R. Stinson
University of Waterloo
Waterloo, ON, Canada N2L 3G1
dstinson@math.uwaterloo.ca

W.D. Wallis
Southern Illinois University-Carbondale
Carbondale, IL, USA 62901-4408
jcmcc@math.siu.edu

On f -colorings of the corona product of cycles with some other graphs

Adiwijaya*, A.N.M. Salman, D. Suprijanto, E.T. Baskoro

Combinatorial Mathematics Research Group
Faculty of Mathematics and Natural Sciences
Institut Teknologi Bandung (ITB)
Jl. Ganesa 10 Bandung 40132

kang_ady@students.itb.ac.id, {msalman, dsuprijanto, ebaskoro}@math.itb.ac.id

Abstract. Let $G = (V(G), E(G))$ be a simple graph and f be a function from $V(G)$ to a subset of positive integers. An f -coloring of G is a generalized edge-coloring such that every vertex $v \in V(G)$ has at most $f(v)$ edges colored with a same color. The minimum number of colors needed to define an f -coloring of G is called an f -chromatic index of G , and denoted by $\chi'_f(G)$. The f -chromatic index of G is equal to $\Delta_f(G)$ or $\Delta_f(G) + 1$, where $\Delta_f(G) = \max\{[d(v)/f(v)] \mid v \in V(G)\}$. G is called in the *class-1*, denoted by C_f1 , if $\chi'_f(G) = \Delta_f(G)$; otherwise G is called in the *class-2*, denoted by C_f2 . In this paper, we showed that the corona product of a cycle with either the complement of a complete graph, or a path, or a cycle is in C_f1 .

Keywords : corona product, edge coloring, f -coloring, f -chromatic index

1 Introduction and statement of results

We consider *simple graphs*, namely finite undirected graphs without loops or multiple edges. Let $G = (V(G), E(G))$ be a graph with the vertex set $V(G)$ and the edge set $E(G)$. Let f be a function from $V(G)$ to a subset of the positive integers. An f -coloring of G is a coloring of edges such that each vertex v has at most $f(v)$ edges colored with a same color. The minimum number of colors needed to define an f -coloring of G is called an f -chromatic index of G , denoted by $\chi'_f(G)$. If $f(v) = 1$ for every $v \in V(G)$, the f -coloring is the classical edge-coloring.

The f -coloring problem is how to determine $\chi'_f(G)$ of a given graph G . It arises in many applications, including the network design problem, the

* **Permanent Address:** Faculty of Science, Institut Teknologi Telkom (IT Telkom), Jl. Telekomunikasi no.1, Dayeuh Kolot Bandung, Indonesia 40257.

scheduling problem, and the file transfer problem in a computer network [1, 2]. The file transfer problem in a computer network is modeled as follows. Each computer is represented by a vertex and every two computers in the file transfer process is represented by an edge. Each computer v has a limit number $f(v)$ of communication ports. If we assume that the transfer time is constant for every file, we can use an f -coloring to manage transferring all files along the minimum time needed.

Let,

$$\Delta_f(G) = \max_{v \in V(G)} \left\{ \left\lceil \frac{d(v)}{f(v)} \right\rceil \right\}. \quad (1)$$

Hakimi and Kariv [3] showed that

$$\Delta_f(G) \leq \chi'_f(G) \leq \Delta_f(G) + 1. \quad (2)$$

G is called in the *class-1*, denoted by C_f1 , if $\chi'_f(G) = \Delta_f(G)$; otherwise G is called in the *class-2*, denoted by C_f2 . Holyer [5] proved that the edge-coloring problem is an NP-complete. It is reduced from the 3SAT problem. Consequently, the f -coloring problem is an NP-complete problem.

Hakimi and Kariv [1] showed that any bipartite graph is in C_f1 . If G is a graph with even $f(v)$ for each $v \in V(G)$, then G is in C_f1 . Yu et al. [7] gave sufficient conditions for fans and wheels to be in C_f1 . Zhang and Liu [9] found the f -chromatic index for complete graphs and gave a classification of complete graphs on f -coloring. In 2008, Zhang et al. [10] presented a classification of regular graphs on f -coloring.

Let G and H be two graphs with n and m vertices, respectively. The *corona product* of a graph G with a graph H , denoted by $G \odot H$, is a graph obtained by taking one copy of an n -vertex graph G and n copies of H , namely H_1, H_2, \dots, H_n , and then for $i = 1, 2, \dots, n$, joining the i -th vertex of G to every vertex of H_i . Here G is called the *center* of $G \odot H$ and H is called the *outer* of $G \odot H$. In this paper, we determine the class of G which obtained by the corona product of a cycle with either the complement of a complete graph, or a path, or a cycle. Let C_m, K_m^c and P_m denote the cycle, the complement of a complete graph and a path on m vertices, respectively. Our results are shown in the following three theorems.

Theorem 1. *Let $n \geq 3$ and $m \geq 1$, $G = C_n \odot K_m^c$, and f be a function from $V(G)$ to a subset of positive integers, then $G \in C_f1$.*

Theorem 2. *Let $n \geq 3$ and $m \geq 2$, $G = C_n \odot P_m$, and f be a function from $V(G)$ to a subset of positive integers, then $G \in C_f1$.*

Theorem 3. *Let $n \geq 3$ and $m \geq 3$, $G = C_n \odot C_m$, and f be a function from $V(G)$ to a subset of positive integers, then $G \in C_f1$.*

Let $C = \{c_1, c_2, \dots, c_{\Delta_f(G)}\}$ be a set of $\Delta_f(G)$ colors. An edge colored with $c \in C$ is called an c -edge. The number of c -edges of G which incident with a vertex v is denoted by $d(v, c)$.

Let,

$$V_0^*(G) = \left\{ v \mid \frac{d(v)}{f(v)} = \Delta_f(G), v \in V(G) \right\}, \quad (3)$$

and

$$V^*(G) = \left\{ v \mid \left\lceil \frac{d(v)}{f(v)} \right\rceil = \Delta_f(G), v \in V(G) \right\}. \quad (4)$$

Zhang and Liu in [8] gave some sufficient conditions for a graph to be in C_f1 as follows.

Lemma 1. [8] (Zhang dan Liu, 2006)

Let G be a graph. If the subgraph induced by $V_0^(G)$ is forest, then $G \in C_f1$.*

Lemma 2. [8] (Zhang dan Liu, 2006)

Let G be a graph. If $f(v^) \nmid d(v^*)$ for every $v^* \in V^*(G)$, then $G \in C_f1$.*

The graphs $G_1 = C_5 \odot K_4^c$, $G_2 = C_5 \odot P_4$, and $G_3 = C_5 \odot C_4$ with $f(v) = 2$ for every v in the center of G and $f(v) = 1$ for every v in the outer of G do not fulfill the premise of Lemma 1 or Lemma 2. But the graphs are in C_f1 as we show in the next section.

2 Proof of theorems

Let $V(C_n^*)$ be the vertex set of the center of G . We label all vertices in $V(C_n^*)$, respectively, by v_1, v_2, \dots, v_n such that $E(C_n^*) = \{v_1v_2, v_2v_3, \dots, v_{n-1}v_n, v_nv_1\}$.

Proof of Theorem 1.

Let,

$$f^* = \min_{v \in V(C_n^*)} \{f(v)\}.$$

Let a multiset $C^* = \{c_i \in C \mid \text{multiplicity of } c_i \text{ is } f^* \text{ for every } i\}$. Now, we color all edges of G by using some colors in C^* . First, we color all edges which are incident with v_1 by using some colors in C^* such that for every $i = 1, 2, \dots, \Delta_f(G)$, the number of c_i -edges which are incident with v_1 is at most f^* . Next, for $j = 2, 3, \dots, n - 1$, we can color respectively, all edges which are incident with v_j by using the colors in $C^* \setminus \{r_j\}$ where r_j is the color that has been used for $v_{j-1}v_j$, respectively. Finally, we color all edges which incident with v_n by using colors in $C^* \setminus \{s, t\}$ where s and t are the colors that have been used for v_1v_n and $v_{n-1}v_n$, respectively. So, all edges of G

have been colored by using $\Delta_f(G)$ colors and every vertex $v \in V(G)$ has at most $f(v)$ edges with a same color. $\square\square$

To prove Theorem 2 and Theorem 3, for $i = 1, 2, \dots, n$, we label all vertices, respectively, in P_m or C_m which adjacent to $v_i \in V(C_n^*)$ by $w_{i,j}$ for $j = 1, 2, \dots, m$. Next, we proof that the corona product of a cycle with a path is in C_f1

Proof of Theorem 2.

It is trivial for case $\Delta_f(G) = 1$. For $\Delta_f(G) \geq 2$, we divide the proof into three cases as follows.

Case 1. $\Delta_f(G) = 2$

In this case, we construct an f -coloring of G as follows. For $i = 1, 2, \dots, n$, $j = 1, 2, \dots, \lceil \frac{m+2}{2} \rceil$, and $\lceil \frac{m+2}{2} \rceil \leq k \leq m - 1$, we color all edges $v_i w_{i,j}$ and $w_{i,k} w_{i,k+1}$ by c_1 . Finally, we color the other edges by c_2 .

Case 2. $\Delta_f(G) = 3$

First, for $i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor$, we color all edges $v_{2i-1} v_{2i}$ by c_1 and for $i = 1, 2, \dots, \lceil \frac{n-2}{2} \rceil$, we color all edges $v_{2i} v_{2i+1}$ by c_2 . Next, for $i = 2, 3, \dots, n - 1$ and $j = 1, \dots, m$, we color all edges $v_i w_{i,j}$ by c_3, c_1, c_2 , alternately. For $i = 1, 2, \dots, n$ and $j = 1, 2, \dots, m - 1$, we color all edges $w_{i,j} w_{i,j+1}$ by c_2, c_3, c_1 , alternately.

Next, we divide the proof into two subcases.

Subcase 2.1. If n is even, for $i = 1$ or $i = n$ and $j = 1, 2, \dots, m$, we color all edges $v_i w_{i,j}$ by c_3, c_1, c_2 , alternately, and the edge $v_n v_1$ by c_2 .

Subcase 2.2. If n is odd, for $j = 1, 2, \dots, m$, we color edges $v_1 w_{1,j}$ by c_2, c_3, c_1 , alternately, and $v_n w_{n,j}$ by c_1, c_2, c_3 , alternately. Finally, we color the edge $v_n v_1$ by c_3 .

Case 3. $\Delta_f(G) \geq 4$

If $v \in V^*$, then $v \in V(C_n^*)$. If $V_0^* = \emptyset$, by Lemma 2, then $G \in C_f1$. If $V_0^* \subset V(C_n^*)$, by Lemma 1, then $G \in C_f1$. If $V_0^* = V(C_n^*)$, we color all edges as follows. For $i = 1, 2, \dots, n$ and $j = 1, 2, \dots, m$, we color $v_i w_{i,j}$ by $c_1, c_2, \dots, c_{\Delta_f(G)}$, alternately. If n is odd, then we replace the color of the first of $c_{\Delta_f(G)}$ -edges which incident with v_n by $c_{\Delta_f(G)-1}$. For $i = 1, 2, \dots, n$ and $j = 1, \dots, m - 1$, we color all edges $w_{i,j} w_{i,j+1}$ by $c_3, c_4, \dots, c_{\Delta_f(G)}, c_1, c_2$, alternately. Finally, for $i = 1, 2, \dots, n - 1$, we color all edges $v_i v_{i+1}$ by $c_{\Delta_f(G)-1}$ and $c_{\Delta_f(G)}$, alternately, and $v_n v_1$ by $c_{\Delta_f(G)}$.

So, all edges of G have been colored by using $\Delta_f(G)$ colors and every vertex $v \in V(G)$ has at most $f(v)$ edges with a same color. $\square\square$

In the next, we prove that the corona product of a cycle with a cycle to be in C_f1 .

Proof of Theorem 3.

For case $\Delta_f(G) = 1$, it is trivial since we can define an f -coloring of G with one color. For $\Delta_f(G) \geq 2$, we divide the proof into four cases as follows.

Case 1. $\Delta_f(G) = 2$

In this case, we construct an f -coloring of G as follows. First, we color all edges in $E(C_n^*)$ by c_1 . For $i = 1, 2, \dots, n$ and $j = 1, 2, \dots, m-1$, we color $v_i w_{i,j}$ by c_2, c_1 , alternately, and $v_i w_{i,m}$ by c_2 . Finally, for $i = 1, 2, \dots, n$ and $j = 1, 2, \dots, m-1$, we color $w_{i,j} w_{i,j+1}$ by c_1, c_2 , alternately, and $w_{i,m} w_{i,1}$ by c_1 .

Case 2. $\Delta_f(G) = 3$

In this case, the proof is divided into three subcases as follows.

Subcase 2.1 $m \equiv 0 \pmod{3}$

First, for $i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor$, we color all edges $v_{2i-1} v_{2i}$ by color c_1 and for $i = 1, 2, \dots, \lceil \frac{n-2}{2} \rceil$, we color all edges $v_{2i} v_{2i+1}$ by c_2 . For even n , we color the last edge $v_n v_1$ by c_2 . For odd n , we color the last edge $v_n v_1$ by c_3 . For $i = 1, 2, \dots, n$ and $j = 1, 2, \dots, m$, we color $v_i w_{i,j}$ by c_1, c_2, c_3 , alternately. Finally, for $i = 1, 2, \dots, n$ and $j = 1, 2, \dots, m-1$, we color all edges $w_{i,j} w_{i,j+1}$ by c_3, c_1, c_2 , alternately.

Subcase 2.2 $m \equiv 1 \pmod{3}$

First, for $i = 1, 2, \dots, n$ and $j = 3, \dots, m-2$, we color $v_i w_{i,j}$ by c_1, c_2, c_3 , alternately. We color $v_i w_{i,1}, v_i w_{i,2}, v_i w_{i,m-1}, v_i w_{i,m}$ for $i = 1, 2, \dots, n$ by c_3, c_3, c_1 , and c_1 , respectively. For $i = 1, 2, \dots, n$ and $k = 1, 2, \dots, m-1$, we color all edges $w_{i,k} w_{i,k+1}$ by c_1, c_2, c_3 , alternately, and $w_{i,m} w_{i,1}$ by c_2 . Finally, we color all edges in $E(C_n^*)$ by c_2 .

Subcase 2.3 $m \equiv 2 \pmod{3}$

For $i = 1, 2, \dots, n$ and $j = 1, 2, \dots, 5$, we color $v_i w_{i,j}$ by c_1, c_2, c_2, c_2 , and c_3 , respectively. For $i = 1, 2, \dots, n$ and $6 \leq j \leq m$, we color edges $v_i w_{i,j}$ by c_1, c_2, c_3 , alternately. Next, for $k = 1, 2, 3$, we color $w_{i,k} w_{i,k+1}$ by c_3, c_1, c_3 , respectively. For $4 \leq k \leq m$, we color $w_{i,k} w_{i,k+1}$ by c_1, c_2, c_3 , alternately, and $w_{i,m} w_{i,1}$ by c_2 . Finally, we color all edges in $E(C_n^*)$ by c_1, c_3 , alternately.

Case 3. $\Delta_f(G) = 4$

If $v \in V^*$, then $v \in V(C_n^*)$. If $V_0^* = \emptyset$, by Lemma 2, then $G \in C_f 1$. If $V_0^* \subset V(C_n^*)$, by Lemma 1, then $G \in C_f 1$. If $V_0^* = V(C_n^*)$, we color all edges as follows. For $i = 1, 2, \dots, n$ and $j = 2, 3, \dots, m-1$, we color $v_i w_{i,j}$ by c_1, c_2, c_3, c_4 , alternately, $v_i w_{i,1}$ by c_1 , and $v_i w_{i,m}$ by c_2 . If n is odd, then we replace the color of the first of c_4 -edges which incident with v_n by c_3 . For $i = 1, 2, \dots, n$ and $j = 1, 2, \dots, m-2$, we color all edges $w_{i,j} w_{i,j+1}$ by c_2, c_3, c_4, c_1 , alternately. Next, for $i = 1, \dots, n$, we color $w_{i,m-1} w_{i,m}$ by c_3 and $w_{i,m} w_{i,1}$ by c_4 . Finally, for $i = 1, 2, \dots, n-1$, we color all edges $v_i v_{i+1}$ by c_3, c_4 , alternately, and $v_n v_1$ by c_4 .

Case 4. $\Delta_f(G) \geq 5$

By the same argument in the Case 3, if $V_0^* = \emptyset$ or $V_0^* \subset V(C_n^*)$, we obtain $G \in C_f 1$. If $V_0^* = V(C_n^*)$, we color all edges as follows. For $i = 1, 2, \dots, n$

and $j = 1, \dots, m$, we color $v_i w_{i,j}$ by $c_1, c_2, \dots, c_{\Delta_f(G)}$, alternately. If n is odd, then we replace the color of the first of $c_{\Delta_f(G)}$ -edges which incident with v_n by $c_{\Delta_f(G)-1}$. For $i = 1, 2, \dots, n$ and $j = 1, 2, \dots, m-1$, we color all edges $w_{i,j} w_{i,j+1}$ by $c_3, c_4, \dots, c_{\Delta_f(G)}, c_1, c_2$, alternately, and the edge $w_{i,m} w_{i,1}$ by $c_{\Delta_f(G)}$. Finally, for $i = 1, 2, \dots, n-1$, we color all edges $v_i v_{i+1}$ by $c_{\Delta_f(G)-1}, c_{\Delta_f(G)}$, alternately, and $v_n v_1$ by $c_{\Delta_f(G)}$.

So, all edges of G have been colored by using $\Delta_f(G)$ colors and every vertex $v \in V(G)$ has at most $f(v)$ edges with a same color. \square

Remark : Base on the proof of Theorem 3 Subcase 2.3, we conclude that $W_m = C_m + \{w\}$ is in $C_f 1$, if $m = 2 \pmod{3}$ and $f(v) \neq 1$ for some $v \in V(C_m)$. So, the conjecture of Yu et al. in [7], namely: a wheel W_m is in $C_f 2$ if $\Delta_f(W_m) = 3$ and $d(w) = 2 \pmod{3}$ with w is core of W_m , is not correct.

References

1. H. Choi, S.L. Hakimi, Scheduling file transfer for trees and odd cycles, *SIAM Journal of Computation* **16:1** (1987) 162-168.
2. E. G. Coffman, M.R. Garey, D.S. Johnson, A.S. LaPaugh, Scheduling file transfers, *SIAM Journal of Computation* **14:3** (1985), 744-780.
3. S.L. Hakimi, O. Kariv, A generalization of edge-coloring in graphs, *Journal of Graph Theory* **10** (1986) 139-154.
4. N. Hartsfield, G. Ringel, *Pearls in Graph Theory : A Comprehensive Introduction*, Academic Press (1994), London.
5. I. Holyer, The NP-completeness of edge-coloring, *SIAM Journal of Computation* **10:4** (1981) 718 - 720.
6. S. Nakano, T. Nishizeki, N. Saito, On the f -coloring of multigraphs, *IEEE Transaction Circuit System* **35:3** (1988) 345-353.
7. J. Yu, L. Han, G. Liu, Some result on the classification for f -colored graphs, *Proceeding ISORA* (2006) .
8. X. Zhang, G. Liu, Some sufficient conditions for a graf to be $C_f 1$, *Applied Mathematics Letters* **19** (2006) 38-44.
9. X. Zhang, G. Liu, The classification of K_n on f -colorings, *Journal of Applied Mathematics and Computing* **19:1-2** (2006) 127-133.
10. X. Zhang, J. Wang, G. Liu, The classification of regular graphs on f -colorings, *Ars Combinatoria* **86** (2008) 273-280.
11. X. Zhou, T. Nishizeki, Edge-coloring and f -coloring for various classes of graphs, *Journal of Graph Algorithm and Applications* **3:1** (1999) 1-18.

On the nonexistence of graphs of diameter 2 and defect 2 by <i>Mirka Miller, Minh Hoang Nguyen and Guillermo Pineda-Villavicencio</i>	5
On Direregularity of Digraphs of Defect at Most Two by <i>Dafik, Mirka Miller, Costas Iliopoulos and Zdenek Ryjacek</i>	21
On distance Magic Labeling of Graphs by <i>K.A. Sugeng, D. Froncek, M. Miller, J. Ryan and J. Walker</i>	39
On the Existence of Regular Supermagic Graphs by <i>Jaroslav Ivanco, Petr Kovar and Andrea Semanicova-Fenovikova</i>	49
An Algorithm for the Optimization of Multiple Classifiers in Data Mining Based on Graphs by <i>Andrei Kelarev, Joe Ryan and John Yearwood</i>	65
TCAM representations of intervals of integers encoded by binary trees by <i>Wojciech Fraczak, Wojciech Rytter and Mohammadreza Yazdani</i>	87
Regular Expression Matching Algorithms using Dual Position Automata by <i>Hiroaki Yamamoto</i>	103
A Compressed Text Index on Secondary Memory by <i>Rodrigo Gonzalez and Gonzalo Navarro</i>	127
Algorithms for Two Versions of LCS Problem for Indeterminate Strings by <i>Costas Iliopoulos, M. Sohel Rahman and Wojciech Rytter</i>	155
On the Ramsey number for a linear forest versus a cocktail party graph by <i>Kashif Ali, A.Q. Baig and Edy Tri Baskoro</i>	173
Vertex-antimagic total labeling of the union of suns by <i>Andrea Parestu, Denny R. Silaban and Kiki A. Sugeng</i>	179
Expansion techniques on the super edge antimagic total graphs by <i>I Wayan Sudarsana, Edy Tri Baskoro, Saladin Uttunggadewa and Dasa Ismailmuza</i>	189
Vertex-magic total labelings of union of generalized Petersen graphs and union of special circulant graphs by <i>Denny R. Silaban, Andrea Parestu, Bong N. Herawati, Kiki A. Sugeng and Slamim</i>	201
The partition dimension of a complete multipartite graph, a special caterpillar and a windmill by <i>Darmaji, S. Uttunggadewa, R. Simanjuntak and E.T. Baskoro</i>	209
Super (a,d)-vertex antimagic total labeling on a disjoint union of regular graphs by <i>Kiki A. Sugeng and Denny R. Silaban</i>	217
On the total vertex-irregular strength of a disjoint union of t copies of a path by <i>Nurdin, A.N.M. Salman, N.N. Gaos and E.T. Baskoro</i>	227
On f -colorings of the corona product of cycles with some other graphs by <i>Adiwijaya, A.N.M. Salman, D. Suprijanto and E.T. Baskoro</i>	235

(Continued on inside back cover)

CONTENTS (continued from back cover)

Notes on exponents of asymmetric two-colored digraphs by Saib Suwilo	243
The Ramsey number of a certain forest respect to a small wheel by I. Wayan Sudarsana, E.T. Baskoro, Hilda Assiyatun and Saladin Uttubggadewa	257
Path-path size multipartite Ramsey numbers by Syafrizal Sy, E.T. Baskoro, S. Uttungadewa and H. Assiyatun	265
On (a,d) - H -antimagic coverings of graphs by Nur Inayah, A.N.M. Salman and R. Simanjuntak	273
The metric dimensions of a complete n -partite graph and its cartesian product with a path by S.W. Saputro, E.T. Baskoro, A.N.M. Salman and D. Suprijanto	283

Subscription Information

All subscription correspondence should be addressed to: Charles Babbage Research Centre, P.O. Box 272, St. Norbert Postal Station, Winnipeg, MB, Canada R3V 1L6. Email: mathjournals@attglobal.net. All back issues of the journal are available.

Information for Authors

Manuscripts of papers should be sent to the Editor-in-Chief, raldred@maths.otago.ac.nz, or to any member of the Editorial Board. It is understood that the contents of the paper have not been, and will not be published elsewhere. Your paper should be prepared in one of the various flavours of TeX (we use AMSLaTeX). The page size should be 4.5 by 7.125 inches (11.4 by 18.1 cm) in 10 point Times or Computer Modern font and the text should be single-spaced, without excessive white spaces.

If you **do** pay page charges, your paper will receive **Priority Publication**. It will go to the head of the queue and appear in the next issue of the journal (this could cut as much as 3 years off the waiting time). This policy will not cause any delay for papers already in the queue, since all priority publication articles will result in additional pages in the journal. Authors who are not able to pay priority publication charges will also benefit considerably, since the priority articles will be removed from the queue and thus the queue will be shortened.

The priority-publication charge is US\$50.00 per page. It is useful if you indicate your intention to pay page charges as early as possible.